

The growth of the Möbius function on the permutation poset

David Marchant

Permutation Patterns 2019

Universität Zürich (UZH), Switzerland.

18th June 2019

The (*principal*) Möbius function $\mu[\pi]$ is defined on the permutation poset as

$$\mu[\pi] = \begin{cases} 1 & \text{if } \pi = 1, \\ -\sum_{\lambda < \pi} \mu[\lambda] & \text{otherwise} \end{cases}$$

The (*principal*) Möbius function $\mu[\pi]$ is defined on the permutation poset as

$$\mu[\pi] = \begin{cases} 1 & \text{if } \pi = 1, \\ -\sum_{\lambda < \pi} \mu[\lambda] & \text{otherwise} \end{cases}$$

Hall's Theorem:

$$\mu[\pi] = \sum_{c \in \mathcal{C}(\pi)} (-1)^{|c|}$$

where $\mathcal{C}(\pi)$ is the set of chains between 1 and π .

What do we mean by the growth of $\mu[\pi]$?

Let $\max_{\mu}(n)$ be the largest absolute value of $\mu[\pi]$ when π is length n .

Then “growth” attempts to place limits on $\max_{\mu}(n)$ as a function of n .

What do we mean by the growth of $\mu[\pi]$?

Let $\max_{\mu}(n)$ be the largest absolute value of $\mu[\pi]$ when π is length n .

Then “growth” attempts to place limits on $\max_{\mu}(n)$ as a function of n .

Previously:

- Burstein, Jelínek, Jelínková and Steingrímsson (2011): Is growth unbounded?
- Smith (2014): Growth is at least $O(n^2)$
- Jelínek, Kantor, Kynčl and Tancer (2018): Growth is at least $O(n^7)$.

What do we mean by the growth of $\mu[\pi]$?

Let $\max_{\mu}(n)$ be the largest absolute value of $\mu[\pi]$ when π is length n .

Then “growth” attempts to place limits on $\max_{\mu}(n)$ as a function of n .

Previously:

- Burstein, Jelínek, Jelínková and Steingrímsson (2011): Is growth unbounded?
- Smith (2014): Growth is at least $O(n^2)$
- Jelínek, Kantor, Kynčl and Tancer (2018): Growth is at least $O(n^7)$.

Today:

- Growth is at least exponential.


What is a 2413-balloon?

What is a 2413-balloon?

	•			
				•
		β		
•				
			•	

With β not the identity permutation $(1, 2, \dots, n)$ or its reverse.











What is a 2413-balloon?



With β not the identity permutation $(1, 2, \dots, n)$ or its reverse.

This is equivalent to the inflation $25314[1, 1, \beta, 1, 1]$.


Reductions of a 2413-balloon



Assume that β is a 2413-balloon, and let π be the 2413-balloon of β

Finding the growth

Assume that β is a 2413-balloon, and let π be the 2413-balloon of β



Partitioning the chains

Partition the chains in the poset $[1, \pi]$:

Partition the chains in the poset $[1, \pi]$:

- $\mathcal{R}$: Chains where the second-highest element is a reduction

Partition the chains in the poset $[1, \pi]$:

- $\mathcal{R}$: Chains where the second-highest element is a reduction
- $\mathcal{G}$: Chains where the largest non-2413-balloon is ≤ 2413 .

Partition the chains in the poset $[1, \pi]$:

- $\mathcal{R}$: Chains where the second-highest element is a reduction
- $\mathcal{G}$: Chains where the largest non-2413-balloon is ≤ 2413 .
- $\mathcal{B}$: All remaining chains.

Partition the chains in the poset $[1, \pi]$:

- $\mathcal{R}$: Chains where the second-highest element is a reduction
- $\mathcal{G}$: Chains where the largest non-2413-balloon is ≤ 2413 .
- $\mathcal{B}$: All remaining chains.

So by Hall's Theorem

$$\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|} + \sum_{c \in \mathcal{G}} (-1)^{|c|} + \sum_{c \in \mathcal{B}} (-1)^{|c|}$$

Can we find an expression for $\sum_{c \in \mathcal{G}} (-1)^{|c|}$?

Can we find an expression for $\sum_{c \in \mathcal{G}} (-1)^{|c|}$?

$$\text{Define } \Phi_G(c) = \begin{cases} \{c\} \cup 2413 & \text{if } 2413 \notin c \\ \{c\} \setminus 2413 & \text{otherwise} \end{cases}$$

Can we find an expression for $\sum_{c \in \mathcal{G}} (-1)^{|c|}$?

$$\text{Define } \Phi_G(c) = \begin{cases} \{c\} \cup 2413 & \text{if } 2413 \notin c \\ \{c\} \setminus 2413 & \text{otherwise} \end{cases}$$

This is an parity-reversing involution on the set $\mathcal{G}$.

Can we find an expression for $\sum_{c \in \mathcal{G}} (-1)^{|c|}$?

$$\text{Define } \Phi_G(c) = \begin{cases} \{c\} \cup 2413 & \text{if } 2413 \notin c \\ \{c\} \setminus 2413 & \text{otherwise} \end{cases}$$

This is an parity-reversing involution on the set $\mathcal{G}$.

$$\text{So } \sum_{c \in \mathcal{G}} (-1)^{|c|} = 0$$

Can we find an expression for $\sum_{c \in \mathcal{G}} (-1)^{|c|}$?

$$\text{Define } \Phi_G(c) = \begin{cases} \{c\} \cup 2413 & \text{if } 2413 \notin c \\ \{c\} \setminus 2413 & \text{otherwise} \end{cases}$$

This is an parity-reversing involution on the set $\mathcal{G}$.

$$\text{So } \sum_{c \in \mathcal{G}} (-1)^{|c|} = 0$$

There is a similar parity reversing involution on the set $\mathcal{B}$.

Can we find an expression for $\sum_{c \in \mathcal{G}} (-1)^{|c|}$?

Define $\Phi_G(c) = \begin{cases} \{c\} \cup 2413 & \text{if } 2413 \notin c \\ \{c\} \setminus 2413 & \text{otherwise} \end{cases}$

This is an parity-reversing involution on the set $\mathcal{G}$.

So $\sum_{c \in \mathcal{G}} (-1)^{|c|} = 0$

There is a similar parity reversing involution on the set $\mathcal{B}$.

So $\sum_{c \in \mathcal{B}} (-1)^{|c|} = 0$

We have $\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|}$. What can we do with this?

We have $\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|}$. What can we do with this?

Partition $\mathcal{R}$ based on the second-highest element. Call this ρ . What do the chains in this sub-partition $(\mathcal{R}_\rho)$ look like?

We have $\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|}$. What can we do with this?

Partition $\mathcal{R}$ based on the second-highest element. Call this ρ . What do the chains in this sub-partition ($\mathcal{R}_\rho$) look like?

Chain in $\mathcal{R}_\rho$




We have $\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|}$. What can we do with this?

Partition $\mathcal{R}$ based on the second-highest element. Call this ρ . What do the chains in this sub-partition $(\mathcal{R}_\rho)$ look like?

Chain in $\mathcal{R}_\rho$



Chain in $[1, \rho]$




We have $\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|}$. What can we do with this?

Partition $\mathcal{R}$ based on the second-highest element. Call this ρ . What do the chains in this sub-partition ($\mathcal{R}_\rho$) look like?

Chain in $\mathcal{R}_\rho$



Chain in $[1, \rho]$



$$\sum_{c \in [1, \rho]} (-1)^{|c|} = \mu[\rho]$$

We have $\mu[\pi] = \sum_{c \in \mathcal{R}} (-1)^{|c|}$. What can we do with this?


Partition $\mathcal{R}$ based on the second-highest element. Call this ρ . What do the chains in this sub-partition ($\mathcal{R}_\rho$) look like?

Chain in $\mathcal{R}_\rho$



$$\sum_{c \in \mathcal{R}_\rho} (-1)^{|c|} = -\mu[\rho]$$

Chain in $[1, \rho]$



$$\sum_{c \in [1, \rho]} (-1)^{|c|} = \mu[\rho]$$

So if **Red** is the set of permutations that are reductions of π , then





$$\begin{aligned}\mu[\pi] &= \sum_{c \in \mathcal{R}} (-1)^{|c|} \\ &= - \sum_{\lambda \in \text{Red}} \mu[\lambda]\end{aligned}$$

Theorem (Burstein, Jelínek, Jelínková and Steingrímsson (2011))

If $\tau \neq 1 \oplus \gamma$, then $\mu[1 \oplus \tau] = -\mu[\tau]$.



Theorem (Burstein, Jelínek, Jelínková and Steingrímsson (2011))

If $\tau \neq 1 \oplus \gamma$, then $\mu[1 \oplus \tau] = -\mu[\tau]$.



Theorem (Burstein, Jelínek, Jelínková and Steingrímsson (2011))



If $\tau \neq 1 \oplus \gamma$, then $\mu[1 \oplus \tau] = -\mu[\tau]$.




$\mu[2\beta 14]$

Theorem (Burstein, Jelínek, Jelínková and Steingrímsson (2011))

If $\tau \neq 1 \oplus \gamma$, then $\mu[1 \oplus \tau] = -\mu[\tau]$.



$\mu[2\beta 14]$





$= -\mu[2\beta 1]$


Handling the reductions

Theorem (Burstein, Jelínek, Jelínková and Steingrímsson (2011))

If $\tau \neq 1 \oplus \gamma$, then $\mu[1 \oplus \tau] = -\mu[\tau]$.



$\mu[2\beta 14]$



$= -\mu[2\beta 1]$





$= \mu[1\beta]$


Handling the reductions

Theorem (Burstein, Jelínek, Jelínková and Steingrímsson (2011))

If $\tau \neq 1 \oplus \gamma$, then $\mu[1 \oplus \tau] = -\mu[\tau]$.



$\mu[2\beta 14]$



$= -\mu[2\beta 1]$



$= \mu[1\beta]$



$= -\mu[\beta]$

Theorem

Let π be a 2413-balloon of β , where β is itself a 2413-balloon. Then $\mu[\pi] = 2\mu[\beta]$.

Theorem

Let π be a 2413-balloon of β , where β is itself a 2413-balloon. Then $\mu[\pi] = 2\mu[\beta]$.

$$\text{Define } \pi^{(n)} = \begin{cases} 1 & \text{If } n = 1 \\ 12 & \text{If } n = 2 \\ 132 & \text{If } n = 3 \\ 2413 & \text{If } n = 4 \\ 2413 \odot \pi^{(n-4)} & \text{Otherwise} \end{cases}$$

Theorem

Let π be a 2413-balloon of β , where β is itself a 2413-balloon. Then $\mu[\pi] = 2\mu[\beta]$.

$$\text{Define } \pi^{(n)} = \begin{cases} 1 & \text{If } n = 1 \\ 12 & \text{If } n = 2 \\ 132 & \text{If } n = 3 \\ 2413 & \text{If } n = 4 \\ 2413 \odot \pi^{(n-4)} & \text{Otherwise} \end{cases}$$

Theorem

For all n , $\max_{\mu}(n) \geq 2^{\lfloor n/4 \rfloor - 1}$.

Future directions...

Can we find a better lower bound for $\max_{\mu}(n)$?

Future directions...

Can we find a better lower bound for $\max_{\mu}(n)$?

What about an upper bound?

We know that $\max_{\mu}(n) \leq n!$. (Zeigler, 1991)

Future directions...

Can we find a better lower bound for $\max_{\mu}(n)$?

What about an upper bound?

We know that $\max_{\mu}(n) \leq n!$. (Zeigler, 1991)

Conjecture

$$\max_{\mu}(n) \leq 2^n$$

Future directions...

Can we find a better lower bound for $\max_{\mu}(n)$?

What about an upper bound?

We know that $\max_{\mu}(n) \leq n!$. (Zeigler, 1991)

Conjecture

$$\max_{\mu}(n) \leq 2^n$$

Make the balloon operation more general. Then what can we say about generalised balloons: $\alpha \odot_{i,j} \beta$?

Future directions...

Can we find a better lower bound for $\max_{\mu}(n)$?

What about an upper bound?

We know that $\max_{\mu}(n) \leq n!$. (Zeigler, 1991)

Conjecture

$$\max_{\mu}(n) \leq 2^n$$

Make the balloon operation more general. Then what can we say about generalised balloons: $\alpha \odot_{i,j} \beta$?

Experimental results up to length 12 indicates that for **most** permutations,

$$\mu[\alpha \odot_{i,j} \beta] \in \{\pm 2\mu[\beta], \pm \mu[\beta], 0\}$$

Future directions...

Can we find a better lower bound for $\max_{\mu}(n)$?

What about an upper bound?

We know that $\max_{\mu}(n) \leq n!$. (Zeigler, 1991)

Conjecture

$$\max_{\mu}(n) \leq 2^n$$

Make the balloon operation more general. Then what can we say about generalised balloons: $\alpha \odot_{i,j} \beta$?

Experimental results up to length 12 indicates that for **most** permutations,

$$\mu[\alpha \odot_{i,j} \beta] \in \{\pm 2\mu[\beta], \pm \mu[\beta], 0\}$$

Further, it appears that $\mu[\alpha \odot_{i,j} \beta] = \pm 2\mu[\beta]$ only if $\beta = \alpha \odot_{i,j} \gamma$.

Thank you!

A. Burstein, V. Jelínek, E. Jelínková and E. Steingrímsson. The Möbius function of separable and decomposable permutations. *Journal of Combinatorial Theory, Series A*, 118(8):2346-2364, 2011.

V. Jelínek, I. Kantor, J. Kynčl, and M. Tancer. On the growth of the Möbius function of permutations. <http://arxiv.org/abs/1809.05774>, 2018.

J. P. Smith. On the Möbius function of permutations with one descent. *Electronic Journal of Combinatorics*, 21(2):Paper 2.11, 19pp., 2014.

G. M. Ziegler. Posets with maximal Möbius function, *Journal of Combinatorial Theory, Series A* 56(2):203-222, 1991.

Talk based on “2413-balloon permutations and the growth of the Möbius function.”
<https://arxiv.org/abs/1812.05064>, 2019.