

# $k$ -partial permutations and the center of the wreath product $\mathcal{S}_k \wr \mathcal{S}_n$ algebra



Omar Tout  
Faculty of Sciences III, Lebanese University



## Part 1: Definitions

### $k$ -partial permutations

If  $i, k \in \mathbb{N}^*$ , the  $k$ -tuple  $p_k(i)$  is the following set of size  $k$  :

$$p_k(i) := \{(i-1)k+1, (i-1)k+2, \dots, ik\}.$$

If  $d = \sqcup_{i=1}^r p_k(a_i)$ , where  $a_i \in \mathbb{N}^*$  for any  $1 \leq i \leq r$ , we define  $\mathcal{B}_d^k$  to be the following group of permutations:

$$\{\omega \in \mathcal{S}_d \mid \forall 1 \leq i \leq r, \exists 1 \leq j \leq r; \omega(p_k(a_i)) = p_k(a_j)\},$$

where  $\mathcal{S}_d$  is the group of permutations of the set  $d$ .

A  $k$ -partial permutation of  $n \in \mathbb{N}$  is a pair  $(d, \omega)$  where  $d \subset [kn]$  is a disjoint union of some  $k$ -tuples and  $\omega \in \mathcal{B}_d^k$ .  $\mathcal{P}_{kn}^k$  will denote the set of all  $k$ -partial permutations of  $n$ .

### Type of a $k$ -partial permutation

A partition  $\rho$  of  $k$  is a decreasing list of positive integers  $(\rho_1, \rho_2, \dots, \rho_r)$  that sum up to  $k$ . Let  $\omega \in \mathcal{B}_d^k$  be written as a product of disjoint cycles and  $\rho$  be a partition of  $k$ . Consider the collection of cycles  $C_1, \dots, C_l$  such that  $C_1$  contains  $\rho_1$  elements of a certain fixed  $k$ -tuple  $p_k(i)$ ,  $C_2$  contains  $\rho_2$  elements  $p_k(i)$ , etc. Now add the part  $m$  to  $\omega(\rho)$  if  $m$  is the number of  $k$ -tuples that form the cycles  $C_1, \dots, C_l$ . We define  $\text{type}(\omega)$  to be the family of partitions  $(\omega(\rho))_{\rho \vdash k}$ .

### Example

Consider the 3-partial permutation  $(d, \omega)$ , where  $d = p_3(1) \cup p_3(2) \cup p_3(4) \cup p_3(6)$  and

$$\omega = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 10 & 11 & 12 & 16 & 17 & 18 \\ 12 & 10 & 11 & 4 & 5 & 6 & 16 & 18 & 17 & 1 & 2 & 3 \end{pmatrix}.$$

Its cycle decomposition is  $(1, 12, 17, 2, 10, 16)(3, 11, 18)(4)(5)(6)$  and its type is formed by  $\omega(2, 1) = (3)$  and  $\omega(1^3) = (1)$ .

## Part 2: The algebra of $k$ -partial permutations

### The algebra of $k$ -partial permutations

The product in  $\mathcal{P}_{kn}^k$  is defined by  $(d_1, \omega_1)(d_2, \omega_2) = (d_1 \cup d_2, \omega_1 \omega_2)$ , where the composition  $\omega_1 \omega_2$  is made after extending both  $\omega_1$  and  $\omega_2$  by identity to  $d_1 \cup d_2$ . Let  $\mathcal{I}_{kn}^k$  be the algebra over  $\mathbb{C}$  generated by the following formal sums indexed by families of partitions  $\Lambda = (\lambda(\rho))_{\rho \vdash k}$  with  $|\Lambda| := \sum_{\rho \vdash k} |\lambda(\rho)| < n$

$$\mathbf{C}_{\Lambda, n} := \sum_{(d, \omega)} (d, \omega),$$

where the sum is taken over all  $(d, \omega) \in \mathcal{P}_{kn}^k$  such that  $|d| = k|\Lambda|$  and  $\text{type}(\omega) = \Lambda$ .

### Homomorphism

There is a surjective homomorphism between the algebras  $\mathcal{I}_{kn}^k$  and  $Z(\mathbb{C}[\mathcal{B}_{kn}^k])$ , the center of the group algebra  $\mathbb{C}[\mathcal{B}_{kn}^k]$ , defined by

$$\psi(\mathbf{C}_{\Lambda, n}) = \binom{n - |\Lambda| + m_1(\Lambda(1^k))}{m_1(\Lambda(1^k))} \mathbf{C}_{\underline{\Lambda}_n},$$

where  $\underline{\Lambda}_n$  is the family of partitions  $\Lambda$  except that  $\Lambda(1^k)$  is replaced by  $\Lambda(1^k) \cup (1^{n-|\Lambda|})$ .

### A universal algebra

Let  $\mathcal{P}_{\infty}^k$  be the group of all the  $k$ -partial permutations with a finite support and  $\mathcal{I}_{\infty}^k$  the algebra generated by the following elements  $\mathbf{C}_{\Lambda} = \sum_{(d, \omega)} (d, \omega)$ , indexed by families of partitions, where the sum runs over all  $(d, \omega) \in \mathcal{P}_{\infty}^k$  such that  $d$  is a union of  $|\Lambda|$   $k$ -tuples and  $\omega$  has type  $\Lambda$ . The projection homomorphism from  $\mathcal{I}_{\infty}^k$  to  $\mathcal{I}_{kn}^k$  is defined by  $\text{Proj}_n(\mathbf{C}_{\Lambda}) = 0$  if  $|\Lambda| > n$  and if  $|\Lambda| \leq n$ ,  $\text{Proj}_n(\mathbf{C}_{\Lambda}) = \mathbf{C}_{\Lambda, n}$ .

## Part 3: Structure coefficients of the center of $\mathcal{B}_{kn}^k$ algebra

### Structure coefficients $c_{\Lambda\Delta}^{\Gamma}$

A family of partitions  $\Lambda$  is called proper if the partition  $\Lambda(1^k)$  does not have any part equal to one. Let  $\Lambda$  and  $\Delta$  be two proper families of partitions with  $|\Lambda|, |\Delta| \leq n$ . In the algebra  $\mathcal{I}_{\infty}^k$ , we can write

$$\mathbf{C}_{\Lambda} \mathbf{C}_{\Delta} = \sum_{\Gamma} c_{\Lambda\Delta}^{\Gamma} \mathbf{C}_{\Gamma},$$

where the sum runs over the families of partitions  $\Gamma$  satisfying  $\max(|\Lambda|, |\Delta|) \leq |\Gamma| \leq |\Lambda| + |\Delta|$  and  $c_{\Lambda\Delta}^{\Gamma}$  are non-negative integers independent of  $n$ . If we apply  $\psi \circ \text{Proj}_n$  to this equality we get the following identity in the center of the group  $\mathcal{B}_{kn}^k$  algebra:

$$\mathbf{C}_{\underline{\Lambda}_n} \mathbf{C}_{\underline{\Delta}_n} = \sum_{\Gamma} c_{\Lambda\Delta}^{\Gamma} \binom{n - |\Gamma| + m_1(\Gamma(1^k))}{m_1(\Gamma(1^k))} \mathbf{C}_{\underline{\Gamma}_n}.$$

The above sum can be turned into a sum over proper families of partitions.

### Polynomiality Property

#### Theorem

Let  $\Lambda, \Delta$  and  $\Gamma$  be three proper families of partitions satisfying  $\max(|\Lambda|, |\Delta|) \leq |\Gamma| \leq |\Lambda| + |\Delta|$ . For any integer  $n \geq |\Gamma|$  we have

$$c_{\underline{\Lambda}_n \underline{\Delta}_n}^{\underline{\Gamma}_n} = \sum_{r=1}^{n-|\Gamma|} c_{\Lambda\Delta}^{\Gamma+r} \binom{n - |\Gamma|}{r},$$

where  $c_{\Lambda\Delta}^{\Gamma+r}$  are non-negative integers independent of  $n$ .

#### Corollary

Let  $\Lambda, \Delta$  and  $\Gamma$  be three proper families of partitions satisfying  $\max(|\Lambda|, |\Delta|) \leq |\Gamma| \leq |\Lambda| + |\Delta|$ . The structure coefficient  $c_{\underline{\Lambda}_n \underline{\Delta}_n}^{\underline{\Gamma}_n}$  is a polynomial in  $n$  with non-negative integer coefficients and of degree at most  $|\Lambda| + |\Delta| - |\Gamma|$ .

## Part 4: An example for $k = 2$

The elements generating  $\mathcal{I}_{\infty}^2$  are indexed by families of partitions  $\Lambda = (\Lambda(1^2), \Lambda(2))$ . For instance,  $C_{((1),(2))}$  is the set of all 2-partial permutations with type  $((1), (2))$  and  $(\{3, 4, 7, 8, 9, 10\}, (3, 7, 4, 8)(9)(10)) \in C_{((1),(2))}$ . In  $\mathcal{I}_{\infty}^2$  :

$$\mathbf{C}_{((1), \emptyset)} \mathbf{C}_{((1),(1))} = 2\mathbf{C}_{((1),(1))} + 2\mathbf{C}_{((1^2), (1))} \text{ and } \mathbf{C}_{(\emptyset, (2))} \mathbf{C}_{(\emptyset, (2))} = 2\mathbf{C}_{((1^2), \emptyset)} + 2\mathbf{C}_{(\emptyset, (1^2))} + 2\mathbf{C}_{(\emptyset, (2^2))} + 3\mathbf{C}_{((3), \emptyset)}.$$

The above red coefficient is due to the fact that there are only two pairs  $(x, y) \in C_{((1), \emptyset)} \times C_{((1),(1))}$  that satisfy  $xy = (\{1, 2, 3, 4\}; (1)(2)(3, 4))$ . Namely  $(x, y)$  can be

$$\left( (\{1, 2\}; (1)(2)), (\{1, 2, 3, 4\}; (1)(2)(3, 4)) \right) \text{ or } \left( (\{3, 4\}; (3)(4)), (\{1, 2, 3, 4\}; (1)(2)(3, 4)) \right).$$

Apply now  $\psi \circ \text{Proj}_n$  for the above two equations to get the following equations in  $Z(\mathbb{C}[\mathcal{B}_{2n}^2])$  :

$$\mathbf{C}_{((1^n), \emptyset)} \mathbf{C}_{((1^{n-1}), (1))} = \mathbf{C}_{((1^{n-1}), (1))} \text{ for any } n \geq 3$$

and for any  $n \geq 5$

$$\mathbf{C}_{((1^{n-2}), (2))} \mathbf{C}_{((1^{n-2}), (2))} = n(n-1)\mathbf{C}_{((1^n), \emptyset)} + 2\mathbf{C}_{((1^{n-2}), (1^2))} + 2\mathbf{C}_{((1^{n-4}), (2^2))} + 3\mathbf{C}_{((1^{n-3}), (3), \emptyset)}.$$

The first equation comes with no surprise since  $C_{((1^n), \emptyset)}$  is the identity class.

### References

- 1- O. Tout, *The center of the wreath product of symmetric groups algebra*, To appear in Algebra and Discrete Mathematics (2019).
- 2- O. Tout, *k-partial permutations and the center of the wreath product  $\mathcal{S}_k \wr \mathcal{S}_n$  algebra*, arXiv e-prints (2019).
- 3- V. Ivanov, S. Kerov, *The algebra of conjugacy classes in symmetric groups, and partial permutations*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) 256(3)(1999) 95–120.