

Classes of Sum Decomposable Affine Permutations

Justin Troyka
York University, Toronto

Joint work with Neal Madras

2019 June 18

Erratum from PP 2018

I neglected to insert a key phrase in my very impressive slide titled *General result on growth rates of sequences*. There, I list several conditions and conclude that, under those conditions,

If $c_n^{(1)} \leq c_n^{(2)}$ for all n and $c_n^{(1)} < c_n^{(2)}$ for some n , then
$$\text{gr}(a_n^{(1)}) < \text{gr}(a_n^{(2)}).$$

It was a very strong theorem.

Erratum from PP 2018

I neglected to insert a key phrase in my very impressive slide titled *General result on growth rates of sequences*. There, I list several conditions and conclude that, under those conditions,

*If $c_n^{(1)} \leq c_n^{(2)}$ for all n and $c_n^{(1)} < c_n^{(2)}$ for some n , then
it is not necessarily the case that $\text{gr}(a_n^{(1)}) < \text{gr}(a_n^{(2)})$.*

It was a very ~~strong~~ **wrong!** theorem.

Affine permutations

- ▶ An **affine permutation** of size n is a bijection $\sigma: \mathbb{Z} \rightarrow \mathbb{Z}$ such that:

(i) $\sigma(i + n) = \sigma(i) + n$ for all $i \in \mathbb{Z}$;

(ii)
$$\sum_{i=1}^n \sigma(i) = \sum_{i=1}^n i.$$

Condition (ii) can also be written as
$$\sum_{i=1}^n (\sigma(i) - i) = 0.$$

- ▶ The affine permutations of size n are an infinite Coxeter group.
- ▶ An affine permutation σ of size n is determined by its values on $\{1, \dots, n\}$, and taking these values mod n results in an (ordinary) permutation.
- ▶ An affine permutation σ of size n is **bounded** if $|\sigma(i) - i| < n$ for all i .

$|\text{AvBA}(321)|$ versus $|\text{AvBA}(312)|$

- ▶ Given a set R of permutations, $\text{AvA}(R)$ is the class of **affine permutations** avoiding every element of R .
 - ▶ For each $n \geq 2$, $\text{AvA}_n(321)$ is infinite.

$|\text{AvBA}(321)|$ versus $|\text{AvBA}(312)|$

- ▶ Given a set R of permutations, $\text{AvA}(R)$ is the class of **affine permutations** avoiding every element of R .
 - ▶ For each $n \geq 2$, $\text{AvA}_n(321)$ is infinite.
- ▶ $\text{AvBA}(R)$ is the class of **bounded affine permutations** avoiding every element of R .
 - ▶ $\text{AvBA}_n(321)$ is finite, but we have not found an exact enumeration.

$|\text{AvBA}(321)|$ versus $|\text{AvBA}(312)|$

- ▶ Given a set R of permutations, $\text{AvA}(R)$ is the class of **affine permutations** avoiding every element of R .
 - ▶ For each $n \geq 2$, $\text{AvA}_n(321)$ is infinite.
- ▶ $\text{AvBA}(R)$ is the class of **bounded affine permutations** avoiding every element of R .
 - ▶ $\text{AvBA}_n(321)$ is finite, but we have not found an exact enumeration.
- ▶ However, $|\text{AvA}_n(312)| = |\text{AvBA}_n(312)| = \frac{1}{2} \binom{2n}{n}$.


I. **Decomposability and the infinite oscillation**

II. Enumeration of decomposable affine permutations

- An affine permutation σ of size n is **sum decomposable** if there is $k \in \mathbb{Z}$ such that

$$i > k \Leftrightarrow \sigma(i) > k.$$


- This is equivalent to the condition that there is an (ordinary) permutation π of size n such that σ has the form $(\cdots \oplus \pi \oplus \pi \oplus \cdots)$, possibly shifted diagonally.



- An affine permutation σ of size n is **sum decomposable** if there is $k \in \mathbb{Z}$ such that

$$i > k \Leftrightarrow \sigma(i) > k.$$

- This is equivalent to the condition that there is an (ordinary) permutation π of size n such that σ has the form $(\cdots \oplus \pi \oplus \pi \oplus \cdots)$, possibly shifted diagonally.




When does an affine permutation class consist only of decomposable elements?

The infinite oscillation

- ▶ The **infinite oscillation** is the affine permutation

$$\mathcal{O} = \dots 30\ 52\ 74\ 96 \dots$$

- ▶ $\mathcal{O}$ has been studied (e.g. by Albert, Brignall, and Vatter) because it can be used to construct **infinite antichains**.




The infinite oscillation

- ▶ The **infinite oscillation** is the affine permutation

$$\mathcal{O} = \dots 30\ 52\ 74\ 96 \dots$$

- ▶ $\mathcal{O}$ has been studied (e.g. by Albert, Brignall, and Vatter) because it can be used to construct **infinite antichains**.



- ▶ $\mathcal{O}$ is indecomposable!

When is an affine permutation class decomposable?

Madras and T. 2019+:

- ▶ An affine permutation is decomposable **if and only if** it avoids $\mathcal{O}$.
- ▶ An affine permutation **class** is decomposable **if and only if** $\mathcal{O}$ is not in it (**if and only if** it is a subset of $\text{AvA}(\mathcal{O})$).
- ▶ $\text{AvA}(\mathcal{R})$ is decomposable **if and only if** there is an element of $\mathcal{R}$ that is contained in $\mathcal{O}$.
 - ▶ Note that the indecomposable permutations contained in $\mathcal{O}$ are the **finite oscillations**

1, 21, 312, 231, 3142, 2413, 31524, 24153, 315264, 241635, ...

I. Decomposability and the infinite oscillation

II. **Enumeration of decomposable affine permutations**

Infinite direct sums

Assume $\mathcal{C}$ is sum closed.

- ▶ Let $\oplus\mathcal{C}$ denote the set of affine permutations obtained by taking infinite direct sums of permutations in $\mathcal{C}$ (possibly with a diagonal shift).
 - ▶ By our theorem, $\oplus\text{Av}(\mathcal{R}) = \text{AvA}(\mathcal{R} \cup \{\emptyset\})$.
- ▶ Let $\mathcal{C}^{\oplus}$ denote the set of (ordinary) permutations in $\mathcal{C}$ that are indecomposable.
- ▶ Madras and T. 2019+: If $A(x)$ and $\tilde{A}(x)$ are the generating functions for $\mathcal{C}$ and $\oplus\mathcal{C}$ respectively, then

$$\tilde{A}(x) = x \frac{d}{dx} \log(A(x)) = \frac{x A'(x)}{A(x)}.$$

- ▶ Madras and T. 2019+: $\max \left\{ |\mathcal{C}_n|, n|\mathcal{C}^\emptyset| \right\} \leq |\oplus \mathcal{C}_n| \leq n|\mathcal{C}_n|$.
 - ▶ In contrast, $|\text{AvAB}_n(321)| \sim (n^2/2)|\text{Av}_n(321)|$.
 - ▶ If an element of $\mathcal{R}$ is contained in $\mathcal{O}$, then this shows that $\text{gr}(\oplus \text{Av}(\mathcal{R})) = \text{gr}(\text{Av}(\mathcal{R}))$ and it answers the question of how much bigger $\oplus \text{Av}_n(\mathcal{R})$ is than $\text{Av}_n(\mathcal{R})$.

- ▶ Madras and T. 2019+: $\max \left\{ |\mathcal{C}_n|, n|\mathcal{C}^\emptyset| \right\} \leq |\oplus \mathcal{C}_n| \leq n|\mathcal{C}_n|$.
 - ▶ In contrast, $|\text{AvAB}_n(321)| \sim (n^2/2)|\text{Av}_n(321)|$.
 - ▶ If an element of R is contained in $\mathcal{O}$, then this shows that $\text{gr}(\oplus \text{Av}(R)) = \text{gr}(\text{Av}(R))$ and it answers the question of how much bigger $\oplus \text{Av}_n(R)$ is than $\text{Av}_n(R)$.

Supercritical and subcritical are technical conditions on the generating function of $\mathcal{C}$ as it relates to the generating function of $\mathcal{C}^\emptyset$. (See Flajolet and Sedgewick, *Analytic Combinatorics*.)

- ▶ Supercritical implies that the **number of blocks** of **almost all** permutations in $\mathcal{C}_n$ is $\sim \alpha n$ for constant $\alpha \in (0, 1)$.
 - ▶ In this situation, $\mathcal{C}^\emptyset$ has a smaller growth rate than $\mathcal{C}$.
 - ▶ Madras and T. 2019+: $|\oplus \mathcal{C}_n| \sim \alpha^{-1} |\mathcal{C}_n| \sim L^n$.
- ▶ Subcritical implies that $|\mathcal{C}_n^\emptyset| \sim \beta |\mathcal{C}_n|$ for some $\beta \in (0, 1)$.
 - ▶ In this situation, $\mathcal{C}^\emptyset$ has the same growth rate as $\mathcal{C}$.
 - ▶ Bóna 2019+: As $n \rightarrow \infty$, $\text{Av}(\pi)$ is subcritical* for almost all $\pi \in S_n^\emptyset$.
 - ▶ Madras and T. 2019+: $|\oplus \mathcal{C}_n| \sim \beta^{1/2} n |\mathcal{C}_n|$.

Examples

R	type	$ Av_n(R) $	$ AvA_n(R) $
312, 231	supercrit.	2^{n-1}	$2^n - 1 \sim 2 Av_n(R) $
312	subcrit.	$\frac{1}{n+1} \binom{2n}{n}$	$\frac{1}{2} \binom{2n}{n} \sim \frac{n}{2} Av_n(R) $
3142, 2413	subcrit.	$\sim k n^{-3/2} \rho^n$	$\sim \frac{n}{\sqrt{2}} Av_n(R) $
3142	subcrit.	$\sim k n^{-5/2} 8^n$	$\sim \frac{27}{32} n Av_n(R) $

$$(\rho = 3 + 2\sqrt{2} \approx 5.828)$$